

Zika Virus Infection

ITV Zika Preparedness


Shelley M. Wood, RN, BSN, CEN

March 16, 2016



Kentucky Public Health
Prevent. Promote. Protect.


Geographic distribution ZIKV 1947-2007



Kentucky Public Health
Prevent. Promote. Protect.

<http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2819875/>

Current Active Zika transmission



Kentucky Public Health
PREVENT. PROMOTE. PROTECT.

WHO declared global health threat of international concern Feb

<http://www.cdc.gov/zika/geo/index.html>

ZIKV in the U.S. 2016

- JAN -Became nationally notifiable condition
- JAN - Imported transmission to the U.S.
- FEB – Sexual transmission in TX
- MAR – First case of Zika in KY



Kentucky Public Health
PREVENT. PROMOTE. PROTECT.

Lab-confirmed Zika in the U.S.A.



11 FEBRUARY

- CONUS
 - Total of 52 travel-associated cases
 - 17 states
- US Territories
 - 173 locally acquired cases
 - 1 travel-associated cases

9 MARCH

- CONUS
 - Total of 193 travel-associated cases
 - 32 states
- US Territories
 - 173 locally acquired cases
 - 1 travel-associated cases



ZIKV Transmission

- Primary route of infection is via bite of infected mosquito



Aedes albopictus



Aedes aegypti

ZIKV Transmission (continued)

Other Modes of Transmission

- Maternal-fetal
 - Intrauterine
 - Perinatal
- Other
 - Sexual
 - Blood transfusion
 - Laboratory exposure
- Theoretical
 - Organ or tissue transplantation
 - Breast milk



Kentucky Public Health
SECURITY • PROMOTE • PROTECT

http://emergency.cdc.gov/coca/ppt/2016/01_z6_16_zika.pdf

Mosquitoes that Carry ZIKV



Aedes aegypti

Aedes species known as container mosquitoes and are aggressive daytime biters

- ***Aedes albopictus***
 - Probable competent vector
 - Known to be infected with Zika
 - Currently, no documentation of transmission
 - Found throughout Kentucky and most states

Aedes albopictus



- ***Aedes aegypti*** mosquito
 - Known competent vector
 - Is common in warm climates including Africa, Asia, South America, Central America, the Caribbean, and the Southeast U.S. and we do have *some* in Kentucky

Clinical Features

- About 20% people infected develop symptoms/
80% remain asymptomatic
- Incubation period
 - 2-12 days
 - Maybe longer
- Characteristic clinical findings: fever, rash, arthralgia, or conjunctivitis.
- Clinical illness is usually mild with symptoms lasting for several days to a week
- Severe disease is uncommon – Hospitalization rare
- Two neurological conditions associated with Zika infection in Brazil and other countries:
 - Microcephaly
 - Guillain-Barre syndrome



Kentucky Public Health
HEALTHY. PROTECTABLE. RESILIENT.



Baby with Typical Head Size



Baby with Microcephaly



Baby with Severe Microcephaly



Recommendations for Testing

- History of travel to Zika-affected area
AND
- Symptomatic within two weeks of travel
- Travel during pregnancy
- Infants born to mom with positive or inconclusive test for Zika virus



Kentucky Testing as of 15 MAR 2016

- Total tests approved = 51
 - 42 females
 - symptomatic
 - 4 pregnant
 - 9 Males
- Thus far, tests resulted from CDC
 - One positive ZIKV



Kentucky Public Health
PROTECT. PROMOTE. PREVENT.

Educate Travelers Prior to U.S. Departure

- Educate all travelers to **wear mosquito repellent for one-two weeks after return from travels**. Best way to prevent Zika virus infection is to avoid mosquito bites by avoiding exposure and eliminating mosquito breeding areas.
- Educate **male travelers to wear condoms** after departure from Zika-affected area for a time period or for the duration of pregnancy of sexual partner
- Until more is known, **pregnant women should consider postponing travel** to any area with active Zika virus transmission
- Healthcare providers should contact their state or local health department about testing patients with compatible travel history and symptoms for Zika
- Educate prevention and control measures
 - Removal and modification of mosquito breeding sites
 - Reducing contact between mosquitoes and people



Kentucky Public Health
PROTECT. PROMOTE. PREVENT.

References

- Zika Virus Outbreak on Yap Island, Federated States of Micronesia
<http://www.nejm.org/doi/pdf/10.1056/NEJMoa0805715>
- World Health Organization Western Pacific Region
http://www.wpro.who.int/mediacentre/factsheets/fs_05182015_zika/en/
- CDC <http://www.cdc.gov/zika/index.html>
- Coca Clinician Outreach and Communication Activity
http://emergency.cdc.gov/coca/ppt/2016/01_26_16_zika.pdf



Kentucky Public Health
PREVENT. PROMOTE. PROTECT.

Questions?

Contact information:

Shelley M. Wood, RN, BSN, CEN
Reportable Diseases Section
Infectious Disease Branch
KY Department for Public Health
shelleym.wood@ky.gov
502.564.3261 EXT. 4242



Kentucky Public Health
PREVENT. PROMOTE. PROTECT.